

## LASIK SURGERY DROP INSTRUCTIONS:



### Zymaxid

- 1 drop in each eye every 2 hours.
- 1 drop in each eye 4 times per day.



### Pred Forte (Use Last)

- 1 drop in each eye every hour.
- 1 drop in each eye every 2 hours.
- 1 drop in each eye 4 times per day.



### Refresh Plus

1 drop in each eye every hour.



### Restasis

1 drop in each eye 2 times per day.

**\*Wait 5 minutes between drops\***

**\*Shake drops well before using\***



## **PRESCRIPTION EYE DROPS AFTER YOUR LASIK PROCEDURE**

- Zymaxid 0.5% will be administered 4 times per day in your treated eye.
- Pred Forte 1% will be administered every 2 hours in your treated eye.
- **DO NOT DISCONTINUE DROPS UNTIL DIRECTED BY DR.HERSCHEL**
- Always wash your hands prior to putting in your eye drops
- The order of drop application is not important
- Remember to refrigerate drops to reduce inflammation
- Use artificial tears as often as needed but at least every 1 to 2 hours. The more you use the better.
- Do not use anything in the treated eye(s) except for medications you have been given by Dr. Herschel (unless otherwise directed by Dr. Herschel).
- To apply your drops simply pull down your lower lid and place the drop in the pocket of the eyelid.
- You may experience a medicinal taste in the back of your throat after putting in your drops, this is normal. You can apply slight pressure with your finger to the side of your nose to help reduce the amount of the medicine that drains into your throat.

**If you have any questions please do not hesitate to contact us at 407-894-8081 between the hours of 9:00 AM to 6:00 PM or you may email us at [info@herschellasik2020.com](mailto:info@herschellasik2020.com)**